

PROTECTIVE ROLE OF ASCORBIC ACID ON CYPERMETHRIN INDUCED ALTERATIONS IN THE PROTEIN LEVELS OF THE FRESHWATER FISH, *PUNTIUS SOPHORE*.

B.R. Shinde.

Jijamata Arts, Science and Commerce College, Nandurbar. 425412.

Abstract

Fresh water Fishes, *Puntius sophore* were exposed to chronic dose of cypermethrin with and without ascorbic acid. Total count of Protein content was recorded. Remarkable decreases in Protein content were observed in cypermethrin exposed fishes. Fishes were exposed to cypermethrin with Ascorbic acid showed less present variation in the Protein content. Pre- exposed fishes to pesticides showed fast recovery with ascorbic acid as compared to cure naturally.

Keyword: cypermethrin, ascorbic acid, Protein, *Puntius sophore*.

Introduction

Indiscriminate use of different pesticides in agriculture to prevent the crop from pest peril has increased over the years, especially in the developing countries (Santhakumar and Balaji, 2000). These pesticides, even when applied in restricted areas are washed and carried away by rains and floods to large water bodies like ponds and rivers and alter the physicochemical properties of water (Waykar and Lomte, 2001), which are proved to be highly toxic not only to fishes but also to other organisms (Madhav Prasad *et.al.*, 2002).

The accumulated pesticide interacts with biomolecules and alters the physiology of organisms. The toxic compounds exert stress to organism and organism responds to it by developing necessary potential to counteract that stress. The chemical changes occurring in the body of organism give first indication of stress (Mayes, 1977). A number of changes in biochemical parameters of aquatic organisms due to pesticide toxicity have been noted by several investigators (Shivaprasad Rao *et. al.* 1981; Mule and Lomte, 1992, Muley *et.al.* 1996). The indiscriminate use of synthetic pyrethroids; organophosphates; organochlorines; carbamates cause serious pollution problem in aquatic environment. Higher concentrations of toxicants in aquatic environment cause adverse effect on aquatic organisms at cellular or molecular level and ultimately lead to alterations in the biochemical composition. The pollutant affects the activity of biologically active molecules such as amino acids, co-enzyme and other proteins containing sulphur and phosphorus, and affect physiological processes in tissues (Ghosh and Chatterjee, 1985) Biochemical changes induced by pesticide are disturbed metabolism, inhibition of important enzymes, retardation of growth and reduction of fecundity and longevity of organisms (Murty, 1986). Reports are available regarding the toxic effects of pesticides on protein content of some aquatic animals (Mule and Lomte , 1993 ; Kaur and Dhawan 1996; Borah and Yadav, 1995). The effect of pesticides are known to induce biochemical changes in the fishes before the drastic cellular dysfunction (Manohar and Subbiah, 1982 and Das and Mukherjee, 2000). Protein is an important organic constituent, which play important role in metabolism of organisms and metabolic activities. It is an integral part of cell membrane. Harper *et.al.* (1978) reported that, the proteins are among the most abundant biological macromolecules and are extremely versatile in their function.

Ascorbic acid is a powerful antioxidant which plays an important role in intracellular oxidation-reduction system and in binding of free radicals produced endogenously (Laurence et.al., 1997). Ascorbic acid reduced the clastogenic effects generated by certain chemical agents in the in vivo and vitro assays. (Amare et.al., 1996, Khan et.al., 1996). Ascorbic acid plays an important role in the distribution and excretion of trace mineral and toxicant Lewin(1974); Hughes(1974) Reported that ascorbic acid is a diffusible biological reducing agent When present in appropriate concentrations. Ascorbic acid acts as an essential factor for normal growth in rainbow trout *salmo gairdneri* reported by (Halver et. al.1969, Tucker and Halver 1986). Chinoy and Seethalakshmi (1977) showed impact of ascorbic acid on steroidogenesis in gastropod. Considering above cited references the effect of pesticides can be detoxified by the ascorbic acid. The present investigation reports the effect of ascorbic acid on cypermethrin induced changes in protein levels of selected tissue of an experimental fresh water fish *puntius sophera*

Materials and Methods

Healthy, active and medium sized fresh water fishes, *puntius sophera* were collected from Shivan river, near Nandurbar. Physicochemical parameters of water used for experimentation were studied by the methods given in APHA and AWWA (1985) The acclimatized, same sized, active healthy fishes were divided into three groups A, B and C. The group A fishes were maintained as control, group B fishes were exposed to chronic dose (LC_{50/10} dose of cypermethrin (0.5910 ppm) for 30days, while group C fishes were exposed to chronic concentration of pesticide with 50 mg/l of ascorbic acid up to 30 days. After 30 days fishes from B group were divided in to two subgroups D and E , the fishes from group D were allowed to self-cure naturally in the normal water , while fishes from E group were expose to 50 mg/l of ascorbic acid Protein content were estimated from A,B and C group fishes after 15th and 30th days of exposure and from D and E groups after 35th and 40th days of recovery, by Lowry's method Lowry's et.al., (1951) using Bovine serum albumin as standard.

Result and Discussion

Biochemical changes in the organs of gill of freshwater fish *Puntius sophera* exposed to chronic concentration of cypermethrin with and without ascorbic acid in Group A,B and C for 15th and 30th days and from Group D and E during recovery after 5 and 10 days were studied along with control fishes with respect to the percentage of protein in dry tissue powders and are given in the table 1.1 the protein contents in the gill of fish in presence of cypermethrin depleted with the increase in exposure period. The changes in the protein contents of a tissue due to pesticide stress indicate the change in the activity of the organism. In presence of ascorbic acid (50 mg/l) the protein depletion is less as compared to those of cypermethrin intoxicated fishes.

The pre-exposed to cypermethrin showed fast recovery in the protein levels in presence of ascorbic acid than those allowed to cure naturally. Proteins are the most important macromolecules in living being which play vital role in architecture and physiology of the cell and in cellular metabolism. Proteins have a major role in the interactions between intra and extra cellular media. As enzymes,Proteins participate in the intricately balanced subcellular function. After the carbohydrate, the next alternative source of energy is protein. To meet the increased energy demand. Decrease in protein may be due to enhance protein catabolism. Cypermethrin causes depletion of protein. The depletion in tissue protein may be due to impairment or low rate of protein synthesis under or due to utilization in the formation of mucoproteins which are released in the form of mucus. Proteins are essential organic constituents which play a vital role in the cellular metabolism. The increase or decrease in the protein contents due to pesticidal stress may be due to changes in the activity of acid phosphatases, rapidchemical binding of enzymes with pesticides, decrease in the synthesis of enzyme protein or increased metabolism due to stress condition (swami et. al.1983).

The decrease in average total protein content in all body tissues after acute and chronic treatment may be due to utilization of amino acids in various metabolic processes (Mule and Lomte,1993), Jha(1988) Supported the idea of consumption of amino acid for metabolic processes as energy source. According to Abel(1974) the depletion of protein may be due to alterations of membrane permeability. Another probability was that pollutant might block protein synthesis. According to Shivaprasad Rao and Ramana Rao(1980) depletion of protein in pollutant treated animal might be due to enhanced proteolytic activity. Increased in protease activity in *Bombax mori* on pesticide treatment also supported depletion of protein content (Srnivas and Purushottam Rao,1987), Waykar(1998) also observed increased protease activity in freshwater bivalve, *Parreysia cylindrica* after pesticide treatment Gopala Rao et. al.(2006) reported that protein serve as energy.

Table No: 1
Physico - chemical parameters of water used for experimentation.

Sr. No.	Parameter	Value
1	Temperature	$25.2 \pm 3.2^{\circ}\text{C}$
2	pH	7.60 ± 0.4
3	Conductivity	$140 \pm 15.7 \mu\text{mho/cm}$
4	Dissolved O ₂	$6.3 \pm 1.1 \text{ ml/l}$
5	Free CO ₂	$3.34 \pm 1.3 \text{ ml/l}$
6	Total Alkalinity	$585.6 \pm 32.8 \text{ mg/l}$
7	Total Hardness	$204 \pm 12.0 \text{ mg/l}$
8	Chloride	$107.92 \pm 16.34 \text{ mg/l}$
9	Calcium	$30.46 \pm 3.06 \text{ mg/l}$
10	Magnesium	$31.67 \pm 2.9 \text{ mg/l}$

Table 1.1: Protein contents in the gill of *Puntius sophore* after chronic exposure to cypermethrin without and with ascorbic acid and during recovery. (Values represent percentage in dry weight).

Group	Treatment	15 days	30 days	35 days	40 days
A	Control	63.65 ± 0.3301	64.48 ± 0.3674	--	--
B	Cypermethrin	$43.48 \pm 0.2442^{***}$ (-31.67)	$47.87 \pm 0.1148^{***}$ (-25.72)	--	--
C	Cypermethrin +AA	$58.68 \pm 0.2262^{***}$ (-7.79)	$55.79 \pm 0.2530^{****}$ (-11.91)	--	--
D	Recovery in Normal water	--	--	$52.43 \pm 0.5228^{###}$ [+9.51]	$58.70 \pm 0.0129^{###}$ [+24.14]
E	Recovery in AA	--	--	$58.65 \pm 0.0140^{###}$ [+22.52]	$62.15 \pm 0.0207^{##}$ [+29.80]

AA = Ascorbic acid (50 mg/l.) \pm Indicates S.D. of three observations. Values in () indicates percent change over respective control. Values in [] indicates percent change over 30 days of respective B.

* indicates significance with the respective control. # indicates significance with 30 days of respective B. $p < 0.05 = * \& ^\#$, $p < 0.01 = ** \& ^\#^*$, $p < 0.001 = *** \& ^\#^*$, NS = Not significant.

References

Abel P.D. (1974): Toxicity of synthetic detergents to fish and aquatic vertebrates. *J. Fish. Biol.* 6: 270-298.

Amare-Mokrane Y.A., Lebucher-Michal M.P., Balavard G., Dumimil G. and Botta B. (1996): Protective effects of L-hederin chlorophyll and ascorbic acid towards the induction of micronuclei by doxorubicin in culture human lymphocytes, mutagensis, and 11:161.

APHA,AWWA and WPCF,(1985): Standard methods for examination of water and wastewater. APHA(17th ed.) Inc. New York.

Borah Sabita and Yadav R.M.S.(1995): Alteration in the protein, free amino acid, nucleic acid and carbohydrate contents of muscle and gill in rogor exposed freshwater fish, *Heteropneustes fossilis*. *Poll. Res.* 14(1): 99-103.

Chinoy N. J. and Seethalakshmi L. (1977): Biochemical evidece for the occurrence of a possible steroid mechanism in reproductive organs of slug, *Laevicaulis alte e*. In Advances. in invertebrates reproduction, Kerala Vol. 1: 365-366.

Das B.K. and Mukherjee S.C. (2000). *Toxicol.* 114 (1-3):11.

Gopala Rao N. Veeraiah K.M., Vijay Kumar and H.Dilleswa Rao, Toxicity and effect of Kelthane (Dicofol 18.5% EC) an organochlorine insecticide to the fresh water fish Channa punctatus, *J. Aqua.Biol.*, 21(2),228-233,(2006).

Ghosh T.K. and Chatterjee S.K. (1985): effect of chromium on tissue energy reserve in freshwater fish, *Sarotherodon mossambicus*. *Environ.Ecol.* 312:178-179.

Halver J.E.(1967): Water soluble Vitamine requirment of chinook Salmon Nutrition of salmonid fishes. *J. Nutr.* 62:225-243.

Hughes R.E.(1974): Nutritional interaction between vitamin C recent aspect of it's physiological and technological importance. Birch G. G. and Parker K. J. (Edition) Applied Science Publisher LTD. London U.K.

Jha B.S. (1988): Effect of lead nitrate on certain organs of air breathing teleost, *Channa punctatus*. Ph.D.Thesis, L.N. Mithila University, Darbhanga (India).

Kaur K. and Dhawan A. (1996): Effect of carbaryl on tissue composition, maturation and breeding potential of *Cirrhina mrigala (Ham.)* bull. *Environ. Contam. Toxicol.* 57:480-486.

Khan P.K. and Sinha S.P. (1996): Effect Ameliorating effect of vitamin C on marine sperm toxicity induced by three pesticides (Endosulfan, phosphamidon and mancozeb) *Mutagenesis*, 11 33.

Lewin(1974): Vitamin C;It's molecular Biology and medical potential. Academic press, London U.K.

Madhav Prasad, Bandyopadhyaya and Ajit Kumar Aditya (2002): Xenobiotic impact on sensitivity in *Anabas testudineus* (Bloch) *J. Ecobiol.* 14(2), 117-124.

Mayes P. A. (1977) : In Review of physiological chemistry 16th edition, Eds Harper H. A. , Rodwell U. V. and Mayes P. A. Lauge medical publications California

Mule M.B. and Lomte, V.S. (1993): Total potein alterations in tissues of freshwater snail, *Thiara tuberculata* during cyperkill exposure Uttar Pradesh. *J. Zool.* 3(1): 63-67.

Muley D.V., Kamble G.B. and Gaikwad P.T. (1996) : Endosulfan toxicity in freshwater fish , *Tilapia mossambica*. *Proc. Acad. Envi. Biol.* 5 (1) : 49-55.

Mule M.B. and Lomte V.S. (1992): Effect of monocrotophos on protein content of fresh water snail, *Thiara tuberculata*. *Asian J. Zool. Science* 1: 25-30.

Mule M.B. and Lomte V.S. (1992): Monocos (Monocrotophos) induced alterations in lipid content of the snail, *Thiara tuberculata*. *J. Ecotoxicol. Envi.Monit.* 2(2):125-128.

Manoharan T. and Subbaiah G. N. (1982): Toxic and Sublethal effects of endosulfan on *Barbus stigma* (Pisces: Cyprinidae) *Proc. Indian Acad. Sci. (Anim. Sci.)*, 91, 523-532

Santhakumar M. and Balaji M. (2000): Acute toxicity of an organophosphorous insecticide monocrotophos and it's effect on behavior of an air breathing fish *Anabas testudineus* (Bloch) *J. Environ. Biol.* 21 (2) 121-123.

Sivaprasad Rao, K. Sathya Prasad, K. Madhu C.H. and Ramana Rao, K. Y. (1981): Effect of methyl parathion on tissue protein and excretory products of the snail, *Pila globosa*. *Nat. Acad. Sci. Letters* 4(8): 337-340.

Swami K. S., Jaganath Rao K.S., Satyavelu K.S. Reddy, K. Srinivasa Moorthy, Lingamoorthy, chetty C. S., and Indira K. (1983): The possible metabolic diversion adapted by the freshwater mussel to counteract the toxic metabolic effects of the selected pesticides. *India J. comp. Anim Physiol.* 1: 95-106.

Tucker B. W. and Halver J. E. (1986): Vit. C metabolism in fish comp. *Pathological Bull.* 18: 1 and 6.

Waykar Bhalchandra B. and Lomte V.S. (2001): Acute toxicity of pesticides carbaryl and endosulfan to freshwater bivalves, *Parreysia cylindrica*. *Poll. Res.* 20(1), 25-29.