

# Comparative Leaf anatomy of some species of *Habenaria* Wild (Orchidaceae).

Soumya Mahabaleshwar Hegde<sup>1\*</sup> and K Krishnaswamy<sup>2</sup>

<sup>1</sup>Department of Botany Kuvempu University Shimoga-577129, Karnataka, India

<sup>2</sup>Department of Botany, Sahyadri Science College, Shimoga, Karnataka, India

**ABSTRACT.** The present study analyzed the characteristics in the leaf of Terrestrial Orchidaceae member i.e different species of *Habenaria* Wild (*H. plantaginea* Lindl., *H. ovalifolia* Wight., *H. multicaudata* Sedgw., *H. crinifera* Lindl., *H. heyneana* Lindl., *H. elwesii* Hook.f., *H. longicorniculata* J.Graham., *H. pelorioides* Par. & Rchb. f. (*Odisha cleistantha* S. Mishra). Anatomical characters are important in identification and ecological adaptations. The anatomical characters have served as a tool for the genus's systematic. The plants were collected at different forest types from Shimoga district. Materials were fixed in FAA solution. Transverse sections were obtained by freehand sections, stained with 1% safranin. Leaf anatomical organization of *Habenaria* species (Stomata, epidermis, mesophyll, vascular bundles, hypodermis.) was observed. There was a distinct variation in the anatomical characters of the leaf. Detailed leaf anatomical characters can be used to distinguish species from others. Leaf anatomical characters with structural adaptation help plants to survive in different vegetation. The implication of this study is the importance of leaf anatomical features to support species identification and to increase understanding of orchid biology which are important in orchid conservation.

**KEYWORDS:** Orchidaceae, Anatomical character, *Habenaria* Wild, Adaptations.

## INTRODUCTION

Orchidaceae with 25,000-30,000 species shows a high level of specialization and a great capacity for adaptation to different environments, which have contributed to the morphological and physiological, vegetative organization, that vary between species (Dressler, 1993; Frander *et al.*, 2017; Udupa, 2011). Orchidaceae is one of the largest family among monocots. Orchidaceae is the second largest flowering plant family of herbaceous, perennials that includes terrestrial, saprophytic, lithophytic, and epiphytic species (Pridgeon *et al.*, 1993). Orchids are associated with an endophytic symbiotic fungus, mainly in their root terrestrial orchids mainly grows in the floor of the forest and grassland.

*Habenaria* Wild is an orchid genus of about 600 species widely distributed throughout the tropical, subtropical and temperate regions of the world. This genus of terrestrial orchids is one of the genera having the largest number of species. In India, it is represented by 17 species in Western Himalaya (Jalal and Jayanthi, 2015). In *Habenaria* Wild. Prominent paired tuberoids with a small third tuberoid is also seen in this species (Pande *et al.*, 2010).

Flowers are highly attractive. The species can be easily identified when in bloom, but the vegetative characteristic (number and size of tubers and leaves, stem height) overlap in many of these. These structural and physiological variations permit plants to survive and reproduce in a variety of environmental conditions and contribute to protection against stress and herbivore damage (Frander *et al.*, 2017.) Particularly in the anatomy of groups are of a descriptive nature, and have focused upon the search for similarities or differences that contribute to the taxonomic determination.

## MATERIALS AND METHODS

### STUDY AREA

Shimoga district is a part of Malnad region of Karnataka and is also known as the 'Gateway to Malnad', There is a seven taluks Soraba, Sagara, Hosanagara, Shimoga, Shikaripura, Thirthalli and Bhadravathi, district situated between the latitudes  $13^{\circ} 27'$  and  $14^{\circ} 39'$  N and between the longitudes  $74^{\circ} 38'$  and  $76^{\circ} 04'$  at a mean altitude of 640 meters above sea level (NRDMS Centre Z. P Office Shimoga). Peak Kodachadri hill is at an altitude of 1343 meters above sea level is the highest point in the district. Shimoga district is rich in diversity of flora and fauna. The district covers evergreen, semi-evergreen, deciduous, scrub forest, and grasslands. (Figure. 1.)



Fig.1. The map shows Shimoga district is the study area

### METHODOLOGY

The present study was undertaken in Shimoga district during the year 2019-2020. The Terrestrial Orchids of eight different *Habenaria* species were collected at different regions from Shimoga district at different vegetation types includes evergreen, deciduous and open forest type. (Table 1.) *Habenaria* species were identified with the help of available manuals and floras (Gamble, 1935; Rao, 1998). To evaluate the leaf anatomy, samples were fixed in FAA 50% for 48 hours and then transferred to Ethanol 50% (Johansen, 1940). Histological slides were prepared by freehand section. Each one was sectioned in the midline, freehand sections were obtained using a razor blade. Transverse sections were subjected to staining with 1% safranin (Kraus and Arduin, 1997). Leaf sections were analyzed in a light microscope. In leaf section number of layers in the epidermis, hypodermis, mesophyll and vascular bundles were observed and noted.

The epidermis of leaves was peeled from the fresh leaves to observe the structure of the stomata. The most important aspects were recorded with a digital camera attached to an Olympus microscope.

### RESULT AND DISCUSSION

The results of this study showed leaf anatomical features of *Habenaria* Wild Species in Shimoga district, comparable data among the species supports species identification and to helps to understand their structures adaption to different habitat in a district. The leaf anatomical structures showed organizations of components forming leaves, consisted of stomata, cuticle, epidermis, mesophyll and vascular bundles, hypodermis in cells are particular in orchid species.

1. *Habenaria heyneana* Lindl.: Leaf transverse section of *H. heyneana* showed a tetracyclic type of stomata, cuticle was thick and smooth, The epidermic was uniseriate with elongated shaped cells, hypodermis was not clear. Mesophyll was homogeneous consisted of 10-12 thin-walled, round-shaped, parenchymatic cell layers, filled with chlorophylls (chlorenchyma) vascular bundles consisted of xylem and phloem, arranged collaterally. There were 4 large and 4-6 small arches arranged in a row with parenchyma cells bundle sheath surrounded the vascular bundles, sclerenchyma cells in vascular bundle were not observed (Figure 2. A, B. Figure 4. I)

2. *Habenaria Plantaginea* Lindl: *H. Plantaginea* anatomical characters showed that stomata were cyclocytic, cuticle was relatively thick and smooth. The epidermis was a single layer composed of elongated shaped cells. Mesophyll was 13 -15 layers, consisted of elongated parenchymatous cells, chlorophylls were present in the parenchymatous cells of the mesophyll. Vascular bundles were arranged within parenchymatous cells of mesophyll layer. Vascular bundles consisted of 7-9 Arches of xylem and phloem. Bundles sheath was thin-walled surrounds the vascular bundles (Figure 2. F, G, H. Figure 4. A)

3. *Habenaria longicorniculata* J.Graham. : *H. longicorniculata* anemocytic and tetracytic stomata configuration, cuticle layer was thick and smooth, epidermic was uniseriate composed of polygonal-shaped cells. Hypodermis was not observed. Mesophyll was 14-15 layers of homogenous polygonal-shaped parenchymatous cells. Vascular bundles consisted of xylem and phloem arranged collaterally. Sclerenchyma cells were not associated with vascular bundles, bundle sheath cells were thin. There were 8-9 arches of xylem and phloem arranged in a row embedded within parenchymatous cells of mesophyll. (Figure 2. I, G Figure .4 B)

4. *Habenaria elwessi* Hook.f : *H. elwessi* had anemocytic stomata configuration and cuticle was thick and smooth, Epidermis is had one layer made up of elongated shaped cells. Hypodermis was absent or not clear. Mesophyll was Homogenous with 15-16 layers consisting of polygonal-shaped parenchymatous cells. Chlorophylls were present in mesophyll cells. Vascular bundles consisted of xylem and phloem arranged collaterally. There were 9-10 xylem and phloem arches arranged in rows embedded in parenchymatous cells. Vascular bundles are surrounded with thin bundle sheath. (Figure 3. A, B, C Figure 4. J).

5. *Habenaria crinifera* Lindl. : *H. crinifera* had tetracyclic stomata configuration with thick and smooth cuticle. Epidermis is single-layered with elongated cells. Hypodermis was not clearly distinguishable. Mesophyll layer is made with Homogenous parenchymatous cells. Mesophyll cells stores starch grains and water. There is a large water storage cells were present. Mesophyll cells with 12-13 layers of parenchymatous cells, they were polygonal in shape. Vascular bundles with 10-11 xylem and phloem arches arranged in rows embedded within parenchymatous cells. (Figure 3. D, E Figure 4. H)

6. *Habenaria ovalifolia* Wight: *H. ovalifolia* had tetracyclic stomata configuration with thick cuticle layer. The epidermis was uniseriate with polygonal-shaped cells. The hypodermis is not clearly seen. Mesophyll was 14-15 layers with homogenous polygonal shaped parenchyma cells. Vascular bundles consisted of xylem and phloem arranged collaterally. Sclerenchyma cells not associated with vascular bundles. Bundle sheath was indistinct. There were 5-6 arches of vascular bundle arranged in a number of rows embedded within parenchymatous cells of mesophyll. (Figure 3. F, G Figure 4. F, G).

7. *Habenaria multicaudata* Sedgw. : *H. multicaudata* leaves has cyclocytic stomata configuration. The cuticle was thick, the epidermis was a single layer with elongated cells. Hypodermis cells were not clear. Mesophyll was 13-14 layers made with homogenous, rounded to elongated shaped parenchyma cells. Chlorophylls were abundantly present in mesophyll cells. Vascular bundles were arranged with 4-5 arches. Sclerenchyma cells were not associated with bundle sheath cells. Bundle sheath cells are thin cleanly visible. (Figure 2 D, E Figure 4.D).

8. *Habenaria pelorioides* E. C. Parish & Rechb.f. (*Odisha cleistantha* S. Mishra): Leaf transverse section of *H. pelorioides* showed the stomata with anemocytic and tetracyclic configuration. The cuticle was thick and smooth in nature. The epidermis was uniseriate with elongated-shaped cells. Hypodermis was absent or not clear. Mesophyll was homogenous consisted of 14-15 layers with thin walled parenchyma cells. Vascular bundles consisted of xylem and phloem arranged collaterally. There were 4-5 arches arranged in rows embedded with parenchyma cells. Sclerenchyma cells were not associated with vascular bundles. Thin-walled bundle sheath surrounded the vascular bundle. (Figure 3. H. I. Figure 4. C, E).

## DISCUSSION

The leaf outline was 'V' shaped in all *Habenaria* species. In all species studied, the cuticle was smooth on the leaf surface. The cuticle thickness was greatest in all studied species. The cuticle is a thick, hydrophobic membrane composed of a polymer matrix (cutin) and associated with solvent-soluble lipids (cuticular waxes) (Rasmussen, 1987); Angela *et al.*, 2015). Plant cuticle plays an important role in the interaction of the plant with the environment, helps to reduce water transpiration (Fahn, 1982; Rindyastuti *et al.*, 2018). *Habenaria* species has a variation in cuticle thickness, the thick cuticle is an ecological adaption to reduce transpiration (Moreira *et al.*, 2013).

The epidermis is the outer cell layer of a leaf, act as a barrier between the leaves and environment it is also important in maintaining the external stimulus (Dietz and Hartung, 1996). The epidermis is important to protect mesophyll and vascular bundles from higher solar radiation. Thick epidermis cell helps to reduce water transpiration.


The large epidermal cell in many species of orchids serves water storage (Guan *et al.*, 2011). Size of the epidermal cell is significantly different among species. Larger epidermal cells support the species to be more adaptive to the warmer environment. The epidermal cell has various shape among species such as polygonal, isodiametric, rectangular and elongated. (Aybake *et al.*, 2010).

The hypodermis is a structure beneath the epidermis this structure can be present or absent in orchid species (Pridgeon, 1982; Stern, 1997; Rindyastuti *et al.*, 2018). Along with epidermis, hypodermis protects the mesophyll cells and vascular bundles.

**Table 1. Ecological traits of *Habenaria* species collected in Shimoga district.**


| Sl.no | SPECIES   | SOURCE                                 | FLOWERING PERIOD    | VEGETATION TYPE  |
|-------|---|--|---------------------|--|
|       |   | locality                               |                     |  |
| 1.    | <i>Habenaria longicorniculata</i> J. Graham.  | Shimoga, hosanagara, Sharavati valley  | August- October     | evergreen, semi-evergreen, grasslands                  |
| 2.    | <i>Habenaria elwesii</i> Hook. f  | Soraba, Sagara kodachadri hills        | July- October       | semi-evergreen forest, grasslands, forest edges,       |
| 3.    | <i>Habenaria heyneana</i> . Lindl.  | Shikaripura, Sagara                    | August- October     | Hills, evergreen to semi evergreen forest.             |
| 4.    | <i>Habenaria crinifera</i> Lindl.   | Thirtahalli, Sagara, Sharavati valley  | July- September     | Grasslands, evergreen to semi- evergreen forest, hills |
| 5.    | <i>Habenaria plantaginea</i> Lindl.   | Sagara Bhadravati, Soraba, Shikaripura | July- September     | Semi-evergreen, Deciduous forest Scrub forest          |
| 6.    | <i>Habenaria ovalifolia</i> . Wight.  | Hosanagara,, Sagara                    | August- October     | Semi-evergreen , Scrub forest                          |
| 7.    | <i>Habenaria multicaudata</i> Sedgw.  | Hosanagara, Thirtahalli,               | July- September     | Evergreen forest                                       |
| 8.    | <i>Habenaria pelorioides</i> Par. & Rchb. f. ( <i>Odisha cleistantha</i> S. Mishra) | Sagara                                 | September- December | Evergreen to semi-evergreen forest                     |

|  |  |  |  |  |
|--|--|--|--|--|
|  |  |  |  |  |
|--|--|--|--|--|




**Figure 2.** Transverse section of leaf **A.** *Habenaria heyneana* Lindl **B.** Mesophyll in *H. heyneana* **C.** Vascular region of *H. heyneana* **D.** T.S of *Habenaria multicaudata* Sedgw. **E.** Vascular region *H. multicaudata* **F** T.S of *Habenaria plantaginea* Lindl. **G.** Vascular region of *H. plantaginea*. **H.** Mesophyll in *H. plantaginea*. **I.** T.S of

*Habenaria longicorniculata* J. Graham. **J.** Epidermal region in *H. longicorniculata*. C-cuticle, UE-upper epidermis, M-mesophyll, PVB-primary vascular bundles. X-xylem, PH-phloem, WSC-water storage cells.



**Figure 3.** Transverse section of leaf **A.** *Habenaria elwessi* Hook.f **B.** Mesophyll in *H. elwessi* **C.** Vascular region of *H. elwessi* **D.** T.S of *Habenaria crinifera* Lindl. **E** Vascular region of *H. crinifera* **F.** T.S of *Habenaria ovalifolia* Wight **G.** Vascular region of *H. ovalifolia* **H.** T.S of *Habenaria pelorioides* Par. & Rchb. f. **I.** Vascular region of *Habenaria pelorioides*. C-cuticle, UE-upper epidermis, M-mesophyll, PVB-primary vascular bundles. X-xylem, PH-phloem, WSC -water storage cells.



**Figure 4.** A. Cyclocytic stomata in *Habenaria plantaginea* Lindl. B. Tetracytic Stomata *Habenaria longicorniculata* J. Graham. C. Tetracytic Stomata *Habenaria pelorioides* Par. & Rchb. f. D. Cyclocytic stomata *Habenaria multicaudata* Sedgw E. Anomocytic stomata in *Habenaria pelorioides* Par. & Rchb. f. F, G. Tetracytic Stomata *Habenaria ovalifolia* Wight. H. Tetracytic Stomata *Habenaria crinifera* Lindl I. Tetracytic Stomata in *Habenaria heyneana* Lindl J. Anomocytic stomata *Habenaria elwessi* Hook.f.

In the present study, hypodermis was absent or not clear in the *Habenaria* species.

The mesophyll is an important structure for leaves that are filled with photosynthesis component (chlorophyll) to assimilate nutrients. The number of mesophyll layers in the present study varies. The thicker leaves support the leaves succulence. The succulence level of leaves related to the parenchymal capacity of mesophylls, to provide water supply for photosynthetic process and leaves cells, turgor mainly in the dry environment (Hsiao, 1973; Lack and Evans, 2001; Metusala *et al.*, 2017). In the present study *Habenaria elwessi* Hook.f has more mesophyll layers than the other *Habenaria* species. In the present study *Habenaria* species shows mesophyll layer is made with homogenous parenchymatic cells.

The vascular bundle is a transport system containing xylem and phloem that are important in the water and nutrient transport (Lack and Evans, 2001; Fahn, 1982).

The vascular bundle's arrangement in the mesophyll of *Habenaria* species has aligned in rows in the centre of mesophyll. Vascular bundles arrangement in a single row was reported in *Habenaria cornuta*, *H. holothrix*, *H. monorrhiza*, *H. occidentalis*, *H. odontoptala*, *H. snowtenii*, *H. vaginatum*. (Stern, 1997) in the present study, sclerenchyma cells were not associated with xylem and phloem. Bundle sheath consisted of cells surrounding the vascular bundle. Bundle sheath cells were made with thin-walled parenchymatic cells. Fibre bundles, spiral thickening, raphide bundles were absent or not clearly visible.

## CONCLUSION

All the species studied possessed both morphological and anatomical features which are most suitable to xeromorphic condition, species are efficient in water use and can adapt to very well even under water stress condition. This study implied the importance of anatomical characters in species identification. Taxonomic knowledge and species identification are important in the conservation, Therefore, *Habenaria* species need more attention in conservation than the more adaptive species, for their future survival.

## ACKNOWLEDGEMENT

I would like to thank Dr Krishnaswamy. K research guide for his encouragement and support given at the start of the study. I am also grateful to my colleagues for cooperation to complete my research work.

## CONFLICT OF INTEREST

The author declares that there are no conflicts of interest.

## REFERENCE

- Aybeke M, Sezik E, Olgun G. 2010 Vegetative anatomy of some Ophrys and Dactylorhiza (Orchidaceae) taxa in Trakya region of Turkey. *Flora* 205:73-89 Cypripedioideae (Orchidaceae). *J. Linn.Soc. (Bot.)*, 59, 309-41.
- Ditz KJ, Hartung W, 1996. The leaf Epidermic: Its Ecophysiological significance. Progress in botany. *fortschritte der botanik*. Volume 57. Springer, berlin. Dressler RL. 1993 Phylogeny and classification of the orchid family. Oregon: *Dreoscorides Press*.
- Fahn A 1982. Plant Anatomy, 3<sup>rd</sup> Ed. *Pergamon press*. Oxford press
- Dietz KJ Hartung W.1996. The Leaf Epidermis: Its Ecophysiological Significance. *Progress In Botany / Fortschritte Der botanic*. Volume 57. Springer, berlin.
- Frander B, Riveron–Giro Anne Damn, Alfredo Garcia–Gonzalez, Lislle- Salis Montero, Osiris Agular- Romeo, Neptali Ramirez, Marcial and Guadalupe Nieto. 2017 Anatomy of the invasive Orchid *Oeceoclades maculate*: Ecological implications. *Botanical journal of the Linnean Society*, 184, 94-112
- Gamble JS. 1935. Flora of Presidency of Madras, Botanical Survey of India, Calcutta, India.
- Jalal JS. And J Jayanthi. 2015 An Annotated Checklist of the Orchids of Western Himalaya, India *Lankesteriana* 15 (1): 7-50
- Johansen DA. 1940 Plant Microtechnique. New York: Mc Graw Hill.
- Kraus JE. and Arduin M. 1997 Manual basico de metodos em morfologia vegetal. Editora da Universidade Federal Rural do Rio de Janeiro, Seropedica
- Kumaraswamy Udupa ES, NR Santhosh, MR Gurudeva, and K Krishnaswamy 2011, Distribution of Ground orchids in grasslands of Chikmagalore district, Karnataka *J. Orchid Soc. India*, 25; 81-85
- Lack AJ Evans De. 2001. Instant notes of Plant Biology. Bios scientific Publishers Limited.Oxford.
- Metusala D, Supriatna J, Nisyawati and Sopandie D. 2017 Comparative leaf and root anatomy of two *Dendrobium* species (Orchidaceae) from different habitat in relation to their potential adaptation to drought. AIP Conference proceedings, 1862
- Moreira ASFP, Filho JPDl, Isaias RMDS. 2013. Structural Adaptations Of Two Sympatric Epiphytic Orchids (Orchidaceae) to a cloudy forest environment in rocky outcrops of Southeast Brazil. *Revista De Biologia Tropical* 61(3): 1053-1065.
- Pridgeon AM. 1993. Systematic Leaf Anatomy Of *Caladenia* (Orchidaceae). *Kew bull*48 (3):533-543.
- Pridgeon AM. Chase P. Crib, and FN. Rasmussen. 1999. *Genera Orchidacearum*. Vol.1 Oxford University Press, Oxford.
- Rao TA. 1998. Conservation of Wild Orchids of Kodgu in the Western Ghats, KAAS, Bangalore, India.
- Ridesti rindyastuti, Siti Nurfadilah, apriyono rahadianoro, Lia hapsari. 2018 leaf anatomical characters of four epiphytic orchids of Sempu Island, East Java, Indonesia: the importance in identification and ecological adaptation. *Biodiversitas*, 19:1906-1918
- Satish Pande, Niranjana S, Vishwasrao, Vivek and Datar, Mandar., 2010, Wild Orchids of Northern Western Ghats. 10.13140/RG.2.1.3586.2887.
- Stern Wl. 1997. Vegetative Anatomy of subtribe Orchidinae (Orchidaceae). *Bot J Linn Soc* 124:121136